
## **Self-Reported Kidney Disease and C8/PFOA**

## Adults (>=18YO) Only

**NOTE:** All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported kidney disease and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

				Self-Reported Kidney Disease		
				No/No Response	Yes	Total
Female	Quartile Group C8	1	Count	8131	1041	9172
			% within Quartile Group C8	88.7%	11.3%	100.0%
			% within Self-Reported Kidney Disease	30.9%	33.0%	31.1%
		2	Count	6534	788	7322
			% within Quartile Group C8	89.2%	10.8%	100.0%
			% within Self-Reported Kidney Disease	24.8%	25.0%	24.8%
		3	Count	6079	653	6732
			% within Quartile Group C8	90.3%	9.7%	100.0%
			% within Self-Reported Kidney Disease	23.1%	20.7%	22.8%
		4	Count	5567	675	6242
			% within Quartile Group C8	89.2%	10.8%	100.0%
			% within Self-Reported Kidney Disease	21.2%	21.4%	21.2%
		Total		26311	3157	29468
			% within Quartile Group C8	89.3%	10.7%	100.0%
			% within Self-Reported Kidney Disease	100.0%	100.0%	100.0%
Male	Quartile Group C8	1	Count	4348	494	4842
			% within Quartile Group C8	89.8%	10.2%	100.0%
			% within Self-Reported Kidney Disease	17.9%	19.9%	18.1%
		2	Count	6240	621	6861
			% within Quartile Group C8	90.9%	9.1%	100.0%
			% within Self-Reported Kidney Disease	25.8%	25.0%	25.7%
		3	Count	6465	663	7128
			% within Quartile Group C8	90.7%	9.3%	100.0%
			% within Self-Reported Kidney Disease	26.7%	26.7%	26.7%
		4	Count	7172	704	7876
			% within Quartile Group C8	91.1%	8.9%	100.0%
			% within Self-Reported Kidney Disease	29.6%	28.4%	29.5%
		Total	Count	24225	2482	26707
			% within Quartile Group C8	90.7%	9.3%	100.0%
			% within Self-Reported Kidney Disease	100.0%	100.0%	100.0%

## Participants with Self-Reported Kidney Disease within C8/PFOA Population Quartile Group



## C8/PFOA for Participants Stratified by Self-Reported Kidney Disease

