
Self-Reported Diagnosis of Fibromyalgia in Adults					
Stratified by Gender and Age					

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of Fibromyalgia	Population in Quartile	Num of Reported	Percent in Quartile
Female	1	30-50	No	3012	2913	96.71%
Female	1	30-50	Yes	3012	99	3.29%
Female	1	51-older	No	2738	2578	94.16%
Female	1	51-older	Yes	2738	160	5.84%
Female	2	30-50	No	3036	2927	96.41%
Female	2	30-50	Yes	3036	109	3.59%
Female	2	51-older	No	2749	2594	94.36%
Female	2	51-older	Yes	2749	155	5.64%
Female	3	30-50	No	3015	2899	96.15%
Female	3	30-50	Yes	3015	116	3.85%
Female	3	51-older	No	2744	2598	94.68%
Female	3	51-older	Yes	2744	146	5.32%
Female	4	30-50	No	3022	2903	96.06%
Female	4	30-50	Yes	3022	119	3.94%
Female	4	51-older	No	2739	2592	94.63%
Female	4	51-older	Yes	2739	147	5.37%
Male	1	30-50	No	2601	2590	99.58%
Male	1	30-50	Yes	2601	11	0.42%
Male	1	51-older	No	2668	2655	99.51%
Male	1	51-older	Yes	2668	13	0.49%
Male	2	30-50	No	2621	2608	99.50%
Male	2	30-50	Yes	2621	13	0.50%
Male	2	51-older	No	2649	2630	99.28%
Male	2	51-older	Yes	2649	19	0.72%
Male	3	30-50	No	2603	2598	99.81%
Male	3	30-50	Yes	2603	5	0.19%
Male	3	51-older	No	2662	2652	99.62%
Male	3	51-older	Yes	2662	10	0.38%
Male	4	30-50	No	2608	2596	99.54%
Male	4	30-50	Yes	2608	12	0.46%
Male	4	51-older	No	2659	2645	99.47%
Male	4	51-older	Yes	2659	14	0.53%


NOTE:All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Fibromyalgia and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of Fibromyalgia in Female Adults Stratified by Age



© 2009 West Virginia University School of Medicine

Self- Reported Diagnosis of Fibromyalgia in Male Adults Stratified by Age



© 2009 West Virginia University School of Medicine