


***Self-Reported Diagnosis of Stroke in Adults
Stratified by Gender and Age***

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of Stroke	Population in Quartile	Num of Reported	Percent in Quartile
Female	1	30-50	No	3012	2975	98.77%
Female	1	30-50	Yes	3012	37	1.23%
Female	1	51-older	No	2738	2628	95.98%
Female	1	51-older	Yes	2738	110	4.02%
Female	2	30-50	No	3036	3018	99.41%
Female	2	30-50	Yes	3036	18	0.59%
Female	2	51-older	No	2749	2629	95.63%
Female	2	51-older	Yes	2749	120	4.37%
Female	3	30-50	No	3015	3001	99.54%
Female	3	30-50	Yes	3015	14	0.46%
Female	3	51-older	No	2744	2641	96.25%
Female	3	51-older	Yes	2744	103	3.75%
Female	4	30-50	No	3022	3002	99.34%
Female	4	30-50	Yes	3022	20	0.66%
Female	4	51-older	No	2739	2623	95.76%
Female	4	51-older	Yes	2739	116	4.24%
Male	1	30-50	No	2601	2578	99.12%
Male	1	30-50	Yes	2601	23	0.88%
Male	1	51-older	No	2668	2539	95.16%
Male	1	51-older	Yes	2668	129	4.84%
Male	2	30-50	No	2621	2604	99.35%
Male	2	30-50	Yes	2621	17	0.65%
Male	2	51-older	No	2649	2534	95.66%
Male	2	51-older	Yes	2649	115	4.34%
Male	3	30-50	No	2603	2581	99.15%
Male	3	30-50	Yes	2603	22	0.85%
Male	3	51-older	No	2662	2558	96.09%
Male	3	51-older	Yes	2662	104	3.91%
Male	4	30-50	No	2608	2590	99.31%
Male	4	30-50	Yes	2608	18	0.69%
Male	4	51-older	No	2659	2559	96.24%
Male	4	51-older	Yes	2659	100	3.76%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Stroke and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of Stroke in Female Adults Stratified by Age



Self- Reported Diagnosis of Stroke in Male Adults Stratified by Age

