
SELF-REPORTED DIAGNOSIS OF SJOREN'S SYNDROME BY GENDER AND AGE
GROUP

				Diagnosed Sjogrens		Total
				NO	YES	
Female	Age Group	0-10	Count	2690	0	2690
			Percent (%)	100.0%	.0%	100.0%
		11-20	Count	4770	0	4770
			Percent (%)	100.0%	.0%	100.0%
		21-30	Count	5407	1	5408
			Percent (%)	100.0%	.0%	100.0%
		31-40	Count	5782	9	5791
			Percent (%)	99.8%	.2%	100.0%
		41-50	Count	6400	13	6413
			Percent (%)	99.8%	.2%	100.0%
		51-60	Count	5231	26	5257
			Percent (%)	99.5%	.5%	100.0%
		61-70	Count	3312	17	3329
			Percent (%)	99.5%	.5%	100.0%
		71-80	Count	1566	5	1571
			Percent (%)	99.7%	.3%	100.0%
		>80	Count	559	0	559
			Percent (%)	100.0%	.0%	100.0%
	Total		Count	35717	71	35788
			Percent (%)	99.8%	.2%	100.0%
Male	Age Group	0-10	Count	2703	0	2703
			Percent (%)	100.0%	.0%	100.0%
		11-20	Count	4961	0	4961
			Percent (%)	100.0%	.0%	100.0%
		21-30	Count	4598	1	4599
			Percent (%)	100.0%	.0%	100.0%
		31-40	Count	4880	0	4880
			Percent (%)	100.0%	.0%	100.0%
		41-50	Count	5674	0	5674
			Percent (%)	100.0%	.0%	100.0%
		51-60	Count	5072	1	5073
			Percent (%)	100.0%	.0%	100.0%
		61-70	Count	3405	0	3405
			Percent (%)	100.0%	.0%	100.0%
		71-80	Count	1597	0	1597
			Percent (%)	100.0%	.0%	100.0%
		>80	Count	350	0	350
			Percent (%)	100.0%	.0%	100.0%
	Total		Count	33240	2	33242
			Percent (%)	100.0%	.0%	100.0%



## SELF-REPORTED SJOGRENS SYNDROME IN FEMALES BY AGE GROUP

©2007 West Virginia University School of Medicine